

Table 14.2.1.1
Summary of Dosing and PK Sampling by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=15)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=3)	
Administered dose(mg/kg/day)						
Baseline Assessment						
N	3	1	3	5	3	15
Mean (SD)	0.10 (0.02)	0.18	0.11 (0.01)	0.19 (0.02)	0.22 (0.02)	0.16 (0.05)
Median (Min, Max)	0.09 (0.09, 0.12)	0.18 (0.18, 0.18)	0.11 (0.10, 0.12)	0.19 (0.17, 0.22)	0.23 (0.20, 0.24)	0.18 (0.09, 0.24)
Final Dosing Visit						
N	3	1	3	4	2	13
Mean (SD)	0.10 (0.02)	0.18	0.11 (0.01)	0.20 (0.02)	0.44 (0.06)	0.19 (0.12)
Median (Min, Max)	0.09 (0.09, 0.12)	0.18 (0.18, 0.18)	0.11 (0.10, 0.12)	0.20 (0.18, 0.22)	0.44 (0.40, 0.48)	0.18 (0.09, 0.48)
Days on study drug						
N	3	1	3	5	3	15
Mean (SD)	13.0 (2.0)	15.0	15.7 (1.2)	16.0 (2.8)	13.7 (1.5)	14.8 (2.2)
Median (Min, Max)	13.0 (11.0, 15.0)	15.0 (15.0, 15.0)	15.0 (15.0, 17.0)	15.0 (14.0, 21.0)	14.0 (12.0, 15.0)	15.0 (11.0, 21.0)
Fresh Plasma Sampling	2 (66.7%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (66.7%)	12 (80.0%)
Urine PK Sampling	2 (66.7%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (66.7%)	12 (80.0%)
Iohexol Collection	2 (66.7%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (66.7%)	12 (80.0%)

Table 14.2.1.2
Summary of Dosing and PK Sampling by Enrollment Group
Safety Population - IC2

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=11)
	0.1 mg/kg/day (N=2)	0.2 mg/kg/day (N=2)	0.1 mg/kg/day (N=5)	0.2 mg/kg/day (N=2)	
Administered dose(mg/kg/day)					
Final Dosing Visit					
N	2	2	5	1	10
Mean (SD)	0.03 (0.01)	0.19 (0.03)	0.10 (0.04)	0.17	0.11 (0.06)
Median (Min, Max)	0.03 (0.03, 0.04)	0.19 (0.16, 0.21)	0.10 (0.03, 0.14)	0.17 (0.17, 0.17)	0.11 (0.03, 0.21)
Type of dosing					
Once a day (QD)	2 (100.0%)	2 (100.0%)	4 (80.0%)	1 (50.0%)	9 (81.8%)
Twice a day (BID)	0	0	1 (20.0%)	0	1 (9.1%)
Days on study drug					
N	2	2	5	2	11
Mean (SD)	26.0 (17.0)	36.0 (9.9)	18.0 (6.5)	26.0 (15.6)	24.2 (11.3)
Median (Min, Max)	26.0 (14.0, 38.0)	36.0 (29.0, 43.0)	15.0 (12.0, 28.0)	26.0 (15.0, 37.0)	21.0 (12.0, 43.0)
Fresh Plasma Sampling	2 (100.0%)	2 (100.0%)	5 (100.0%)	1 (50.0%)	10 (90.9%)
Urine PK Sampling	2 (100.0%)	2 (100.0%)	5 (100.0%)	1 (50.0%)	10 (90.9%)
Iohexol Collection	2 (100.0%)	1 (50.0%)	5 (100.0%)	1 (50.0%)	9 (81.8%)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Systolic BP at Baseline						
N	3	1	3	5	2	14
Mean (SD)	119.7 (5.5)	133.0	122.7 (0.6)	125.8 (9.2)	124.5 (9.2)	124.1 (7.0)
Median (Min, Max)	117.0 (116.0, 126.0)	133.0 (133.0, 133.0)	123.0 (122.0, 123.0)	126.0 (114.0, 137.0)	124.5 (118.0, 131.0)	123.0 (114.0, 137.0)
Systolic BP at Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	114.7 (11.1)	127.0	116.0 (4.6)	117.0 (5.5)	113.5 (6.4)	116.5 (6.7)
Median (Min, Max)	116.0 (103.0, 125.0)	127.0 (127.0, 127.0)	117.0 (111.0, 120.0)	119.0 (110.0, 124.0)	113.5 (109.0, 118.0)	117.5 (103.0, 127.0)
Systolic BP at 10 Mins Post-lohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	112.3 (15.9)	131.0	108.0 (8.0)	112.8 (5.9)	97.5 (6.4)	110.6 (11.5)
Median (Min, Max)	120.0 (94.0, 123.0)	131.0 (131.0, 131.0)	108.0 (100.0, 116.0)	113.0 (106.0, 119.0)	97.5 (93.0, 102.0)	110.0 (93.0, 131.0)
Systolic BP at 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	117.0 (2.0)	120.0	110.7 (8.1)	111.0 (8.8)	94.0 (12.7)	110.4 (10.4)
Median (Min, Max)	117.0 (115.0, 119.0)	120.0 (120.0, 120.0)	112.0 (102.0, 118.0)	109.5 (102.0, 123.0)	94.0 (85.0, 103.0)	112.0 (85.0, 123.0)
Systolic BP at 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	115.3 (10.2)	121.0	108.0 (6.2)	106.5 (6.0)	109.5 (10.6)	110.5 (8.0)
Median (Min, Max)	111.0 (108.0, 127.0)	121.0 (121.0, 121.0)	110.0 (101.0, 113.0)	106.0 (100.0, 114.0)	109.5 (102.0, 117.0)	110.0 (100.0, 127.0)
Systolic BP at 12 Hour						
N	3	1	3	4	2	13

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	125.7 (5.5)	126.0	109.0 (1.0)	117.0 (10.5)	106.5 (13.4)	116.2 (10.3)
Median (Min, Max)	123.0 (122.0, 132.0)	126.0 (126.0, 126.0)	109.0 (108.0, 110.0)	119.0 (104.0, 126.0)	106.5 (97.0, 116.0)	116.0 (97.0, 132.0)
Systolic BP at 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	117.3 (4.5)	109.0	120.0 (6.2)	113.8 (9.7)	115.5 (4.9)	115.9 (6.8)
Median (Min, Max)	117.0 (113.0, 122.0)	109.0 (109.0, 109.0)	122.0 (113.0, 125.0)	112.0 (104.0, 127.0)	115.5 (112.0, 119.0)	114.0 (104.0, 127.0)
Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-5.0 (11.4)	-6.0	-6.7 (4.0)	-8.8 (11.7)	-11.0 (2.8)	-7.6 (8.3)
Median (Min, Max)	-10.0 (-13.0, 8.0)	-6.0 (-6.0, -6.0)	-6.0 (-11.0, -3.0)	-10.0 (-19.0, 10.0)	-11.0 (-13.0, -9.0)	-9.5 (-19.0, 10.0)
Change from Baseline to 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-7.3 (14.0)	-2.0	-14.7 (8.0)	-16.0 (5.2)	-27.0 (2.8)	-14.3 (10.2)
Median (Min, Max)	-6.0 (-22.0, 6.0)	-2.0 (-2.0, -2.0)	-14.0 (-23.0, -7.0)	-16.0 (-22.0, -10.0)	-27.0 (-29.0, -25.0)	-14.0 (-29.0, 6.0)
Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-2.7 (3.8)	-13.0	-12.0 (8.2)	-17.8 (14.5)	-30.5 (3.5)	-14.5 (12.3)
Median (Min, Max)	-1.0 (-7.0, 0.0)	-13.0 (-13.0, -13.0)	-10.0 (-21.0, -5.0)	-16.5 (-35.0, -3.0)	-30.5 (-33.0, -28.0)	-10.0 (-35.0, 0.0)
Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-4.3 (14.0)	-12.0	-14.7 (6.4)	-22.3 (11.5)	-15.0 (19.8)	-14.5 (12.3)
Median (Min, Max)	-5.0 (-18.0, 10.0)	-12.0 (-12.0, -12.0)	-12.0 (-22.0, -10.0)	-25.0 (-33.0, -6.0)	-15.0 (-29.0, -1.0)	-12.0 (-33.0, 10.0)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	6.0 (10.0)	-7.0	-13.7 (0.6)	-11.8 (11.9)	-18.0 (4.2)	-8.7 (11.5)
Median (Min, Max)	6.0 (-4.0, 16.0)	-7.0 (-7.0, -7.0)	-14.0 (-14.0, -13.0)	-9.5 (-28.0, 0.0)	-18.0 (-21.0, -15.0)	-12.0 (-28.0, 16.0)
Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-2.3 (9.5)	-24.0	-2.7 (6.7)	-15.0 (11.7)	-9.0 (4.2)	-9.0 (10.5)
Median (Min, Max)	1.0 (-13.0, 5.0)	-24.0 (-24.0, -24.0)	-1.0 (-10.0, 3.0)	-17.0 (-27.0, 1.0)	-9.0 (-12.0, -6.0)	-10.0 (-27.0, 5.0)
Percentage Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-4.1 (9.6)	-4.5	-5.4 (3.3)	-6.5 (9.3)	-8.8 (1.6)	-6.0 (6.7)
Median (Min, Max)	-7.9 (-11.2, 6.8)	-4.5 (-4.5, -4.5)	-4.9 (-9.0, -2.4)	-8.3 (-14.4, 8.8)	-8.8 (-9.9, -7.6)	-7.8 (-14.4, 8.8)
Percentage Change from Baseline to 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-6.2 (12.1)	-1.5	-12.0 (6.5)	-12.4 (3.6)	-21.7 (0.7)	-11.4 (8.2)
Median (Min, Max)	-4.8 (-19.0, 5.1)	-1.5 (-1.5, -1.5)	-11.5 (-18.7, -5.7)	-12.4 (-16.7, -7.9)	-21.7 (-22.1, -21.2)	-11.7 (-22.1, 5.1)
Percentage Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-2.1 (3.0)	-9.8	-9.8 (6.6)	-13.4 (10.4)	-24.7 (4.7)	-11.4 (9.5)
Median (Min, Max)	-0.9 (-5.6, 0.0)	-9.8 (-9.8, -9.8)	-8.2 (-17.1, -4.1)	-12.8 (-25.5, -2.4)	-24.7 (-28.0, -21.4)	-8.2 (-28.0, 0.0)
Percentage Change from Baseline to 8 Hour						
N	3	1	3	4	2	13

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	-3.4 (11.4)	-9.0	-12.0 (5.2)	-17.0 (8.3)	-11.5 (15.1)	-11.2 (9.5)
Median (Min, Max)	-4.3 (-14.3, 8.5)	-9.0 (-9.0, -9.0)	-9.8 (-17.9, -8.1)	-19.4 (-24.1, -5.0)	-11.5 (-22.1, -0.8)	-9.8 (-24.1, 8.5)
Percentage Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	5.3 (8.5)	-5.3	-11.1 (0.5)	-9.0 (8.9)	-14.6 (4.5)	-6.8 (9.3)
Median (Min, Max)	5.1 (-3.2, 13.8)	-5.3 (-5.3, -5.3)	-11.4 (-11.5, -10.6)	-7.3 (-21.2, 0.0)	-14.6 (-17.8, -11.5)	-8.8 (-21.2, 13.8)
Percentage Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-1.7 (7.6)	-18.1	-2.2 (5.4)	-11.5 (8.7)	-7.1 (2.9)	-6.9 (8.0)
Median (Min, Max)	0.9 (-10.3, 4.3)	-18.1 (-18.0, -18.0)	-0.8 (-8.1, 2.5)	-13.5 (-19.7, 0.8)	-7.1 (-9.2, -5.1)	-8.1 (-19.7, 4.3)
Systolic z score at Baseline						
N	3	1	3	5	2	14
Mean (SD)	1.1 (0.3)	1.9	1.5 (0.6)	1.8 (0.7)	2.5 (0.3)	1.7 (0.6)
Median (Min, Max)	1.0 (0.9, 1.4)	1.9 (1.9, 1.9)	1.3 (1.0, 2.1)	1.9 (1.0, 2.7)	2.5 (2.3, 2.7)	1.7 (0.9, 2.7)
Systolic z score at Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	0.6 (0.9)	1.3	0.8 (0.8)	0.9 (0.7)	1.5 (0.0)	0.9 (0.7)
Median (Min, Max)	0.5 (-0.2, 1.6)	1.3 (1.3, 1.3)	0.4 (0.2, 1.8)	0.9 (0.1, 1.9)	1.5 (1.5, 1.5)	1.0 (-0.2, 1.9)
Systolic z score at 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	0.4 (1.3)	1.7	0.1 (0.3)	0.4 (0.5)	-0.0 (0.0)	0.4 (0.8)
Median (Min, Max)	0.9 (-1.1, 1.5)	1.7 (1.7, 1.7)	-0.1 (-0.1, 0.3)	0.5 (-0.1, 0.9)	-0.0 (-0.1, -0.0)	0.2 (-1.1, 1.7)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Systolic z score at 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.9 (0.1)	0.7	0.3 (0.2)	0.3 (0.9)	-0.4 (0.5)	0.4 (0.6)
Median (Min, Max)	0.9 (0.8, 0.9)	0.7 (0.7, 0.7)	0.3 (0.1, 0.5)	0.3 (-0.7, 1.2)	-0.4 (-0.7, 0.0)	0.5 (-0.7, 1.2)
Systolic z score at 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.7 (1.1)	0.8	0.1 (0.1)	-0.2 (1.2)	1.1 (1.6)	0.4 (1.0)
Median (Min, Max)	0.6 (-0.3, 1.8)	0.8 (0.8, 0.8)	0.1 (-0.0, 0.1)	-0.4 (-1.3, 1.5)	1.1 (-0.1, 2.2)	0.1 (-1.3, 2.2)
Systolic z score at 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	1.7 (0.8)	1.2	0.2 (0.6)	0.8 (1.0)	0.8 (0.6)	0.9 (0.9)
Median (Min, Max)	1.5 (1.1, 2.5)	1.2 (1.2, 1.2)	-0.1 (-0.3, 0.8)	1.3 (-0.7, 1.5)	0.8 (0.4, 1.3)	1.1 (-0.7, 2.5)
Systolic z score at 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.9 (0.6)	-0.4	1.2 (0.3)	0.5 (0.5)	1.7 (0.2)	0.9 (0.7)
Median (Min, Max)	1.1 (0.2, 1.4)	-0.4 (-0.4, -0.4)	1.1 (0.9, 1.6)	0.4 (0.1, 1.2)	1.7 (1.6, 1.8)	1.1 (-0.4, 1.8)
Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-0.5 (1.1)	-0.6	-0.6 (0.4)	-0.8 (1.1)	-1.0 (0.3)	-0.7 (0.8)
Median (Min, Max)	-1.0 (-1.2, 0.7)	-0.6 (-0.6, -0.6)	-0.6 (-1.1, -0.3)	-1.0 (-1.8, 0.9)	-1.0 (-1.2, -0.8)	-0.9 (-1.8, 0.9)
Change from Baseline to 10 Mins Post-lohexol Administration						
N	3	1	3	4	2	13

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	-0.7 (1.3)	-0.2	-1.4 (0.8)	-1.5 (0.5)	-2.6 (0.3)	-1.4 (1.0)
Median (Min, Max)	-0.6 (-2.1, 0.6)	-0.2 (-0.2, -0.2)	-1.3 (-2.2, -0.7)	-1.5 (-2.1, -0.9)	-2.6 (-2.8, -2.3)	-1.3 (-2.8, 0.6)
Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.3 (0.4)	-1.2	-1.2 (0.8)	-1.7 (1.4)	-2.9 (0.3)	-1.4 (1.2)
Median (Min, Max)	-0.1 (-0.7, -0.0)	-1.2 (-1.2, -1.2)	-1.0 (-2.0, -0.5)	-1.6 (-3.3, -0.3)	-2.9 (-3.1, -2.7)	-1.0 (-3.3, -0.0)
Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.4 (1.3)	-1.1	-1.4 (0.6)	-2.1 (1.1)	-1.4 (1.9)	-1.4 (1.2)
Median (Min, Max)	-0.5 (-1.7, 0.9)	-1.1 (-1.1, -1.1)	-1.2 (-2.1, -0.9)	-2.3 (-3.2, -0.6)	-1.4 (-2.8, -0.1)	-1.2 (-3.2, 0.9)
Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.6 (1.0)	-0.7	-1.3 (0.0)	-1.1 (1.1)	-1.7 (0.4)	-0.8 (1.1)
Median (Min, Max)	0.6 (-0.4, 1.5)	-0.7 (-0.7, -0.7)	-1.3 (-1.3, -1.2)	-0.9 (-2.6, -0.0)	-1.7 (-2.0, -1.4)	-1.2 (-2.6, 1.5)
Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.2 (0.9)	-2.3	-0.3 (0.6)	-1.4 (1.1)	-0.9 (0.4)	-0.9 (1.0)
Median (Min, Max)	0.1 (-1.2, 0.5)	-2.3 (-2.2, -2.2)	-0.1 (-1.0, 0.3)	-1.6 (-2.6, 0.1)	-0.9 (-1.1, -0.6)	-1.0 (-2.6, 0.5)
Percentage Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-35.4 (104.5)	-30.0	-50.8 (34.2)	-33.0 (74.6)	-41.1 (7.0)	-38.3 (60.2)
Median (Min, Max)	-66.1 (-121.2, 80.9)	-30.0 (-30.0, -30.0)	-56.6 (-81.7, -14.1)	-58.3 (-92.9, 96.7)	-41.1 (-46.0, -36.1)	-51.3 (-121.2, 96.7)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Percentage Change from Baseline to 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-61.3 (133.9)	-10.2	-91.9 (22.6)	-79.8 (20.4)	-101.2 (1.7)	-76.3 (61.4)
Median (Min, Max)	-39.7 (-204.8, 60.4)	-10.2 (-10.2, -10.2)	-104.0 (-105.9, -65.8)	-73.8 (-107.5, -64.2)	-101.2 (-102.4, -100.1)	-83.0 (-204.8, 60.4)
Percentage Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-19.0 (24.0)	-64.5	-72.8 (24.8)	-77.4 (51.3)	-115.5 (23.4)	-67.7 (43.7)
Median (Min, Max)	-9.7 (-46.3, -1.1)	-64.5 (-64.5, -64.5)	-74.3 (-96.8, -47.3)	-79.4 (-125.4, -25.4)	-115.5 (-132.0, -98.9)	-64.5 (-132.0, -1.1)
Percentage Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-21.4 (112.3)	-59.6	-94.7 (6.2)	-119.5 (76.2)	-53.3 (69.5)	-76.3 (74.4)
Median (Min, Max)	-46.9 (-118.8, 101.5)	-59.6 (-59.6, -59.6)	-93.6 (-101.4, -89.1)	-117.8 (-214.5, -28.0)	-53.3 (-102.4, -4.1)	-93.6 (-214.5, 101.5)
Percentage Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	60.7 (87.3)	-34.9	-98.2 (35.7)	-53.3 (58.5)	-68.6 (21.9)	-38.3 (77.2)
Median (Min, Max)	60.4 (-26.5, 148.2)	-34.9 (-34.9, -34.9)	-104.0 (-130.7, -60.0)	-37.8 (-136.7, -0.8)	-68.6 (-84.1, -53.1)	-43.1 (-136.7, 148.2)
Percentage Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-8.9 (69.7)	-118.7	-11.4 (34.2)	-62.7 (47.7)	-33.3 (13.0)	-38.2 (52.2)
Median (Min, Max)	8.8 (-85.8, 50.2)	-118.7 (-118.7, -118.7)	-10.2 (-46.3, 22.2)	-80.6 (-96.8, 7.5)	-33.3 (-42.5, -24.1)	-42.5 (-118.7, 50.2)
Normal (<90th Percentile) Systolic BP at Baseline						
No	1 (33.3%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	2 (100.0%)	9 (64.3%)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Yes	2 (66.7%)	0	1 (33.3%)	2 (40.0%)	0	5 (35.7%)
Normal (<90th Percentile) Systolic BP at Pre-Dose at PK Visit						
No	1 (33.3%)	1 (100.0%)	1 (33.3%)	1 (20.0%)	2 (100.0%)	6 (42.9%)
Yes	2 (66.7%)	0	2 (66.7%)	4 (80.0%)	0	8 (57.1%)
Normal (<90th Percentile) Systolic BP at 10 Mins Post-Iohexol Administration						
No	1 (33.3%)	1 (100.0%)	0	0	0	2 (14.3%)
Yes	2 (66.7%)	0	3 (100.0%)	4 (80.0%)	2 (100.0%)	11 (78.6%)
Normal (<90th Percentile) Systolic BP at 4 Hour						
No	0	0	0	0	0	0
Yes	3 (100.0%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (100.0%)	13 (92.9%)
Normal (<90th Percentile) Systolic BP at 8 Hour						
No	1 (33.3%)	0	0	1 (20.0%)	1 (50.0%)	3 (21.4%)
Yes	2 (66.7%)	1 (100.0%)	3 (100.0%)	3 (60.0%)	1 (50.0%)	10 (71.4%)
Normal (<90th Percentile) Systolic BP at 12 Hour						
No	2 (66.7%)	0	0	2 (40.0%)	0	4 (28.6%)
Yes	1 (33.3%)	1 (100.0%)	3 (100.0%)	2 (40.0%)	2 (100.0%)	9 (64.3%)
Normal (<90th Percentile) Systolic BP at 24 Hour						
No	1 (33.3%)	0	1 (33.3%)	0	2 (100.0%)	4 (28.6%)
Yes	2 (66.7%)	1 (100.0%)	2 (66.7%)	4 (80.0%)	0	9 (64.3%)
Improvement (to <90th Percentile) from Baseline at Pre-Dose at PK Visit						
No	2 (66.7%)	1 (100.0%)	2 (66.7%)	2 (40.0%)	2 (100.0%)	9 (64.3%)
Yes	1 (33.3%)	0	1 (33.3%)	3 (60.0%)	0	5 (35.7%)

Table 14.2.2.1.1.1
Summary of Systolic BP by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Improvement (to <90th Percentile) from Baseline at 10 Mins Post-lohexol Administration						
No	2 (66.7%)	1 (100.0%)	1 (33.3%)	1 (20.0%)	0	5 (35.7%)
Yes	1 (33.3%)	0	2 (66.7%)	3 (60.0%)	2 (100.0%)	8 (57.1%)
Improvement (to <90th Percentile) from Baseline at 4 Hour						
No	2 (66.7%)	0	1 (33.3%)	1 (20.0%)	0	4 (28.6%)
Yes	1 (33.3%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	2 (100.0%)	9 (64.3%)
Improvement (to <90th Percentile) from Baseline at 8 Hour						
No	2 (66.7%)	0	1 (33.3%)	2 (40.0%)	1 (50.0%)	6 (42.9%)
Yes	1 (33.3%)	1 (100.0%)	2 (66.7%)	2 (40.0%)	1 (50.0%)	7 (50.0%)
Improvement (to <90th Percentile) from Baseline at 12 Hour						
No	2 (66.7%)	0	1 (33.3%)	3 (60.0%)	0	6 (42.9%)
Yes	1 (33.3%)	1 (100.0%)	2 (66.7%)	1 (20.0%)	2 (100.0%)	7 (50.0%)
Improvement (to <90th Percentile) from Baseline at 24 Hour						
No	2 (66.7%)	0	2 (66.7%)	1 (20.0%)	2 (100.0%)	7 (50.0%)
Yes	1 (33.3%)	1 (100.0%)	1 (33.3%)	3 (60.0%)	0	6 (42.9%)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Diastolic BP at Baseline						
N	3	1	3	5	2	14
Mean (SD)	76.0 (18.7)	73.0	75.3 (7.8)	75.4 (8.1)	76.5 (16.3)	75.5 (10.2)
Median (Min, Max)	82.0 (55.0, 91.0)	73.0 (73.0, 73.0)	73.0 (69.0, 84.0)	78.0 (63.0, 83.0)	76.5 (65.0, 88.0)	75.5 (55.0, 91.0)
Diastolic BP at Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	72.0 (3.5)	80.0	66.3 (8.6)	69.4 (13.5)	69.5 (14.8)	70.1 (9.9)
Median (Min, Max)	74.0 (68.0, 74.0)	80.0 (80.0, 80.0)	68.0 (57.0, 74.0)	69.0 (50.0, 86.0)	69.5 (59.0, 80.0)	71.5 (50.0, 86.0)
Diastolic BP at 10 Mins Post-lohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	63.3 (10.0)	75.0	74.7 (3.1)	59.5 (12.1)	60.0 (9.9)	65.2 (10.5)
Median (Min, Max)	64.0 (53.0, 73.0)	75.0 (75.0, 75.0)	74.0 (72.0, 78.0)	63.0 (42.0, 70.0)	60.0 (53.0, 67.0)	67.0 (42.0, 78.0)
Diastolic BP at 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	65.0 (4.6)	60.0	67.3 (6.5)	58.8 (6.8)	61.5 (13.4)	62.7 (7.1)
Median (Min, Max)	66.0 (60.0, 69.0)	60.0 (60.0, 60.0)	67.0 (61.0, 74.0)	57.5 (52.0, 68.0)	61.5 (52.0, 71.0)	61.0 (52.0, 74.0)
Diastolic BP at 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	70.0 (10.4)	68.0	75.0 (11.8)	50.5 (4.7)	64.5 (3.5)	64.2 (12.2)
Median (Min, Max)	76.0 (58.0, 76.0)	68.0 (68.0, 68.0)	72.0 (65.0, 88.0)	51.5 (44.0, 55.0)	64.5 (62.0, 67.0)	65.0 (44.0, 88.0)
Diastolic BP at 12 Hour						
N	3	1	3	4	2	13

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	73.7 (2.9)	56.0	70.0 (8.2)	66.8 (12.0)	69.0 (24.0)	68.6 (10.9)
Median (Min, Max)	72.0 (72.0, 77.0)	56.0 (56.0, 56.0)	68.0 (63.0, 79.0)	67.5 (54.0, 78.0)	69.0 (52.0, 86.0)	72.0 (52.0, 86.0)
Diastolic BP at 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	69.7 (8.6)	54.0	69.7 (0.6)	60.3 (9.8)	74.0 (12.7)	66.2 (9.5)
Median (Min, Max)	68.0 (62.0, 79.0)	54.0 (54.0, 54.0)	70.0 (69.0, 70.0)	58.0 (52.0, 73.0)	74.0 (65.0, 83.0)	68.0 (52.0, 83.0)
Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-4.0 (20.0)	7.0	-9.0 (7.5)	-6.0 (6.7)	-7.0 (1.4)	-5.4 (10.0)
Median (Min, Max)	-14.0 (-17.0, 19.0)	7.0 (7.0, 7.0)	-10.0 (-16.0, -1.0)	-7.0 (-13.0, 5.0)	-7.0 (-8.0, -6.0)	-7.5 (-17.0, 19.0)
Change from Baseline to 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-12.7 (19.6)	2.0	-0.7 (9.5)	-14.5 (7.0)	-16.5 (6.4)	-9.9 (12.0)
Median (Min, Max)	-18.0 (-29.0, 9.0)	2.0 (2.0, 2.0)	-1.0 (-10.0, 9.0)	-14.5 (-21.0, -8.0)	-16.5 (-21.0, -12.0)	-10.0 (-29.0, 9.0)
Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-11.0 (15.1)	-13.0	-8.0 (13.1)	-15.3 (8.1)	-15.0 (2.8)	-12.4 (9.6)
Median (Min, Max)	-13.0 (-25.0, 5.0)	-13.0 (-13.0, -13.0)	-2.0 (-23.0, 1.0)	-15.0 (-24.0, -7.0)	-15.0 (-17.0, -13.0)	-13.0 (-25.0, 5.0)
Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-6.0 (27.0)	-5.0	-0.3 (6.7)	-23.5 (9.9)	-12.0 (12.7)	-10.9 (16.0)
Median (Min, Max)	-6.0 (-33.0, 21.0)	-5.0 (-5.0, -5.0)	3.0 (-8.0, 4.0)	-24.5 (-34.0, -11.0)	-12.0 (-21.0, -3.0)	-8.0 (-34.0, 21.0)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	-2.3 (18.1)	-17.0	-5.3 (13.6)	-7.3 (4.8)	-7.5 (7.8)	-6.5 (10.5)
Median (Min, Max)	-5.0 (-19.0, 17.0)	-17.0 (-17.0, -17.0)	-10.0 (-16.0, 10.0)	-7.0 (-13.0, -2.0)	-7.5 (-13.0, -2.0)	-9.0 (-19.0, 17.0)
Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-6.3 (21.2)	-19.0	-5.7 (8.3)	-13.8 (7.1)	-2.5 (3.5)	-8.9 (11.3)
Median (Min, Max)	-3.0 (-29.0, 13.0)	-19.0 (-19.0, -19.0)	-3.0 (-15.0, 1.0)	-15.0 (-20.0, -5.0)	-2.5 (-5.0, 0.0)	-5.0 (-29.0, 13.0)
Percentage Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-0.4 (30.3)	9.6	-11.8 (10.2)	-8.6 (9.7)	-9.2 (0.1)	-6.3 (15.0)
Median (Min, Max)	-17.1 (-18.7, 34.5)	9.6 (9.6, 9.6)	-11.9 (-21.9, -1.4)	-9.7 (-20.6, 6.2)	-9.2 (-9.2, -9.1)	-9.5 (-21.9, 34.5)
Percentage Change from Baseline to 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-12.9 (26.5)	2.7	-0.1 (12.5)	-20.1 (10.7)	-21.2 (3.8)	-12.2 (16.2)
Median (Min, Max)	-19.8 (-35.4, 16.4)	2.7 (2.7, 2.7)	-1.4 (-11.9, 13.0)	-18.3 (-33.3, -10.3)	-21.2 (-23.9, -18.5)	-12.5 (-35.4, 16.4)
Percentage Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-11.4 (18.7)	-17.8	-9.6 (15.5)	-20.2 (9.5)	-19.7 (0.5)	-15.5 (12.0)
Median (Min, Max)	-15.9 (-27.5, 9.1)	-17.8 (-17.8, -17.8)	-2.9 (-27.4, 1.4)	-20.3 (-28.9, -11.1)	-19.7 (-20.0, -19.3)	-17.8 (-28.9, 9.1)
Percentage Change from Baseline to 8 Hour						
N	3	1	3	4	2	13

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	-1.8 (37.5)	-6.9	-0.6 (9.0)	-31.0 (10.8)	-14.2 (13.6)	-12.8 (21.8)
Median (Min, Max)	-7.3 (-36.3, 38.2)	-6.9 (-6.8, -6.8)	4.4 (-11.0, 4.8)	-31.5 (-43.6, -17.5)	-14.2 (-23.9, -4.6)	-11.0 (-43.6, 38.2)
Percentage Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	1.3 (26.7)	-23.3	-6.1 (18.0)	-10.2 (7.2)	-11.1 (12.5)	-7.8 (15.6)
Median (Min, Max)	-6.1 (-20.9, 30.9)	-23.3 (-23.3, -23.3)	-13.7 (-19.0, 14.5)	-10.2 (-18.1, -2.6)	-11.1 (-20.0, -2.3)	-13.7 (-23.3, 30.9)
Percentage Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-4.0 (27.8)	-26.0	-6.8 (9.9)	-18.6 (9.0)	-2.8 (4.0)	-10.7 (15.2)
Median (Min, Max)	-3.7 (-31.9, 23.6)	-26.0 (-26.0, -26.0)	-4.1 (-17.9, 1.4)	-20.8 (-26.4, -6.4)	-2.8 (-5.7, 0.0)	-6.4 (-31.9, 23.6)
Diastolic z score at Baseline						
N	3	1	3	5	2	14
Mean (SD)	1.2 (1.5)	0.8	1.1 (0.9)	1.1 (0.8)	1.6 (1.1)	1.2 (0.9)
Median (Min, Max)	1.8 (-0.5, 2.3)	0.8 (0.8, 0.8)	0.8 (0.4, 2.2)	1.1 (-0.0, 2.1)	1.6 (0.8, 2.4)	1.0 (-0.5, 2.4)
Diastolic z score at Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	0.8 (0.3)	1.4	0.3 (0.9)	0.6 (1.3)	1.0 (1.0)	0.7 (0.9)
Median (Min, Max)	0.8 (0.5, 1.1)	1.4 (1.4, 1.4)	0.3 (-0.6, 1.2)	0.4 (-1.1, 2.2)	1.0 (0.3, 1.7)	0.6 (-1.1, 2.2)
Diastolic z score at 10 Mins Post-Iohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	0.0 (0.8)	1.0	1.1 (0.3)	-0.3 (1.1)	0.2 (0.5)	0.3 (0.9)
Median (Min, Max)	0.2 (-0.9, 0.7)	1.0 (1.0, 1.0)	1.2 (0.7, 1.2)	0.1 (-1.8, 0.4)	0.2 (-0.2, 0.5)	0.5 (-1.8, 1.2)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Diastolic z score at 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.2 (0.4)	-0.3	0.4 (0.5)	-0.4 (0.7)	0.3 (0.8)	0.0 (0.6)
Median (Min, Max)	0.0 (-0.1, 0.6)	-0.3 (-0.3, -0.3)	0.2 (0.1, 0.9)	-0.3 (-1.2, 0.3)	0.3 (-0.3, 0.9)	0.0 (-1.2, 0.9)
Diastolic z score at 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.6 (1.1)	0.4	1.1 (1.3)	-1.1 (0.6)	0.6 (0.0)	0.2 (1.2)
Median (Min, Max)	1.2 (-0.7, 1.3)	0.4 (0.4, 0.4)	0.7 (0.1, 2.5)	-1.1 (-1.8, -0.3)	0.6 (0.5, 0.6)	0.4 (-1.8, 2.5)
Diastolic z score at 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.9 (0.4)	-0.7	0.6 (0.7)	0.3 (1.2)	1.0 (1.8)	0.6 (1.0)
Median (Min, Max)	0.9 (0.6, 1.3)	-0.7 (-0.7, -0.7)	0.7 (-0.0, 1.3)	0.2 (-0.8, 1.7)	1.0 (-0.3, 2.3)	0.7 (-0.8, 2.3)
Diastolic z score at 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	0.6 (0.9)	-0.8	0.6 (0.2)	-0.2 (0.9)	1.4 (0.8)	0.4 (0.9)
Median (Min, Max)	0.6 (-0.3, 1.5)	-0.8 (-0.8, -0.8)	0.6 (0.5, 0.8)	-0.3 (-1.1, 0.7)	1.4 (0.8, 2.0)	0.6 (-1.1, 2.0)
Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-0.4 (1.8)	0.6	-0.8 (0.7)	-0.5 (0.6)	-0.6 (0.1)	-0.5 (0.9)
Median (Min, Max)	-1.3 (-1.6, 1.6)	0.6 (0.6, 0.6)	-0.9 (-1.4, -0.1)	-0.6 (-1.1, 0.4)	-0.6 (-0.7, -0.5)	-0.7 (-1.6, 1.6)
Change from Baseline to 10 Mins Post-lohexol Administration						
N	3	1	3	4	2	13

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Mean (SD)	-1.2 (1.8)	0.2	-0.1 (0.9)	-1.3 (0.6)	-1.5 (0.6)	-0.9 (1.1)
Median (Min, Max)	-1.6 (-2.6, 0.8)	0.2 (0.2, 0.2)	-0.1 (-0.9, 0.8)	-1.3 (-1.8, -0.7)	-1.5 (-1.9, -1.0)	-0.9 (-2.6, 0.8)
Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-1.0 (1.4)	-1.1	-0.7 (1.2)	-1.3 (0.7)	-1.3 (0.3)	-1.1 (0.9)
Median (Min, Max)	-1.2 (-2.3, 0.4)	-1.1 (-1.1, -1.1)	-0.2 (-2.1, 0.1)	-1.4 (-2.1, -0.6)	-1.3 (-1.6, -1.1)	-1.1 (-2.3, 0.4)
Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.6 (2.4)	-0.4	-0.0 (0.6)	-2.1 (0.8)	-1.1 (1.2)	-1.0 (1.4)
Median (Min, Max)	-0.6 (-3.0, 1.8)	-0.4 (-0.4, -0.4)	0.3 (-0.7, 0.4)	-2.2 (-2.9, -1.0)	-1.1 (-1.9, -0.3)	-0.7 (-3.0, 1.8)
Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.2 (1.6)	-1.5	-0.5 (1.2)	-0.7 (0.4)	-0.7 (0.7)	-0.6 (0.9)
Median (Min, Max)	-0.5 (-1.7, 1.5)	-1.5 (-1.5, -1.5)	-0.9 (-1.5, 0.9)	-0.6 (-1.2, -0.2)	-0.7 (-1.1, -0.2)	-0.8 (-1.7, 1.5)
Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-0.6 (1.9)	-1.6	-0.5 (0.8)	-1.2 (0.6)	-0.2 (0.3)	-0.8 (1.0)
Median (Min, Max)	-0.3 (-2.6, 1.1)	-1.6 (-1.6, -1.6)	-0.3 (-1.4, 0.1)	-1.3 (-1.7, -0.4)	-0.2 (-0.5, -0.0)	-0.5 (-2.6, 1.1)
Percentage Change from Baseline to Pre-Dose at PK Visit						
N	3	1	3	5	2	14
Mean (SD)	-149.2 (136.9)	74.8	-78.7 (78.7)	1376.4 (3171.0)	-46.1 (22.8)	441.5 (1903.6)
Median (Min, Max)	-72.9 (-307.2, -67.4)	74.8 (74.8, 74.8)	-42.5 (-168.9, -24.6)	-25.0 (-97.2, 7048.3)	-46.1 (-62.2, -29.9)	-52.4 (-307.2, 7048.3)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Percentage Change from Baseline to 10 Mins Post-lohexol Administration						
N	3	1	3	4	2	13
Mean (SD)	-122.4 (44.3)	21.1	55.5 (143.3)	2773.8 (5726.7)	-101.2 (32.2)	824.1 (3168.2)
Median (Min, Max)	-145.2 (-150.7, -71.4)	21.1 (21.0, 21.0)	-11.0 (-42.5, 219.9)	-71.8 (-124.9, 11364)	-101.2 (-123.9, -78.4)	-71.4 (-150.7, 11364)
Percentage Change from Baseline to 4 Hour						
N	3	1	3	4	2	13
Mean (SD)	-82.4 (15.8)	-140.2	-45.5 (53.9)	839.8 (1983.3)	-98.9 (50.0)	202.9 (1086.3)
Median (Min, Max)	-80.4 (-99.2, -67.7)	-140.2 (-140.1, -140.1)	-49.0 (-97.5, 10.1)	-87.6 (-277.2, 3811.7)	-98.9 (-134.2, -63.5)	-80.4 (-277.2, 3811.7)
Percentage Change from Baseline to 8 Hour						
N	3	1	3	4	2	13
Mean (SD)	-167.3 (157.3)	-54.2	1.8 (80.0)	1326.5 (3096.2)	-54.9 (33.3)	357.4 (1690.5)
Median (Min, Max)	-130.9 (-339.6, -31.3)	-54.2 (-54.2, -54.2)	16.8 (-84.7, 73.2)	-186.2 (-291.1, 5969.4)	-54.9 (-78.4, -31.4)	-78.4 (-339.6, 5969.4)
Percentage Change from Baseline to 12 Hour						
N	3	1	3	4	2	13
Mean (SD)	-125.4 (131.7)	-183.1	23.6 (192.1)	1168.4 (2482.6)	-70.9 (89.5)	311.0 (1381.5)
Median (Min, Max)	-75.4 (-274.8, -26.1)	-183.1 (-183.1, -183.1)	-67.9 (-105.8, 244.3)	-18.3 (-180.3, 4890.6)	-70.9 (-134.2, -7.6)	-67.9 (-274.8, 4890.6)
Percentage Change from Baseline to 24 Hour						
N	3	1	3	4	2	13
Mean (SD)	-113.6 (97.1)	-204.6	-23.8 (44.6)	1396.3 (3050.3)	-9.6 (12.9)	380.7 (1681.6)
Median (Min, Max)	-115.0 (-210.0, -15.7)	-204.6 (-204.6, -204.6)	-32.1 (-63.7, 24.3)	-60.5 (-263.4, 5969.4)	-9.6 (-18.8, -0.5)	-38.3 (-263.4, 5969.4)
Normal (<90th Percentile) Diastolic BP at Baseline						
No	2 (66.7%)	0	1 (33.3%)	2 (40.0%)	1 (50.0%)	6 (42.9%)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Yes	1 (33.3%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	1 (50.0%)	8 (57.1%)
Normal (<90th Percentile) Diastolic BP at Pre-Dose at PK Visit						
No	0	1 (100.0%)	0	2 (40.0%)	1 (50.0%)	4 (28.6%)
Yes	3 (100.0%)	0	3 (100.0%)	3 (60.0%)	1 (50.0%)	10 (71.4%)
Normal (<90th Percentile) Diastolic BP at 10 Mins Post-Iohexol Administration						
No	0	0	0	0	0	0
Yes	3 (100.0%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (100.0%)	13 (92.9%)
Normal (<90th Percentile) Diastolic BP at 4 Hour						
No	0	0	0	0	0	0
Yes	3 (100.0%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	2 (100.0%)	13 (92.9%)
Normal (<90th Percentile) Diastolic BP at 8 Hour						
No	0	0	1 (33.3%)	0	0	1 (7.1%)
Yes	3 (100.0%)	1 (100.0%)	2 (66.7%)	4 (80.0%)	2 (100.0%)	12 (85.7%)
Normal (<90th Percentile) Diastolic BP at 12 Hour						
No	1 (33.3%)	0	1 (33.3%)	1 (20.0%)	1 (50.0%)	4 (28.6%)
Yes	2 (66.7%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	1 (50.0%)	9 (64.3%)
Normal (<90th Percentile) Diastolic BP at 24 Hour						
No	1 (33.3%)	0	0	0	1 (50.0%)	2 (14.3%)
Yes	2 (66.7%)	1 (100.0%)	3 (100.0%)	4 (80.0%)	1 (50.0%)	11 (78.6%)
Improvement (to <90th Percentile) from Baseline at Pre-Dose at PK Visit						
No	1 (33.3%)	1 (100.0%)	2 (66.7%)	5 (100.0%)	2 (100.0%)	11 (78.6%)
Yes	2 (66.7%)	0	1 (33.3%)	0	0	3 (21.4%)

Table 14.2.2.1.1.2
Summary of Diastolic BP by Enrollment Group
Safety Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²			Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	0.4 mg/kg/day (N=2)	
Improvement (to <90th Percentile) from Baseline at 10 Mins Post-Iohexol Administration						
No	1 (33.3%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	1 (50.0%)	8 (57.1%)
Yes	2 (66.7%)	0	1 (33.3%)	1 (20.0%)	1 (50.0%)	5 (35.7%)
Improvement (to <90th Percentile) from Baseline at 4 Hour						
No	1 (33.3%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	1 (50.0%)	8 (57.1%)
Yes	2 (66.7%)	0	1 (33.3%)	1 (20.0%)	1 (50.0%)	5 (35.7%)
Improvement (to <90th Percentile) from Baseline at 8 Hour						
No	1 (33.3%)	1 (100.0%)	3 (100.0%)	3 (60.0%)	1 (50.0%)	9 (64.3%)
Yes	2 (66.7%)	0	0	1 (20.0%)	1 (50.0%)	4 (28.6%)
Improvement (to <90th Percentile) from Baseline at 12 Hour						
No	2 (66.7%)	1 (100.0%)	2 (66.7%)	4 (80.0%)	2 (100.0%)	11 (78.6%)
Yes	1 (33.3%)	0	1 (33.3%)	0	0	2 (14.3%)
Improvement (to <90th Percentile) from Baseline at 24 Hour						
No	2 (66.7%)	1 (100.0%)	2 (66.7%)	3 (60.0%)	2 (100.0%)	10 (71.4%)
Yes	1 (33.3%)	0	1 (33.3%)	1 (20.0%)	0	3 (21.4%)

Table 14.2.2.1.2.1
Summary of Systolic Blood Pressure by Enrollment Group
PD Population - IC2 with Valid Pre-Lisinopril Records

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=11)
	0.1 mg/kg/day (N=2)	0.2 mg/kg/day (N=2)	0.1 mg/kg/day (N=5)	0.2 mg/kg/day (N=2)	
Systolic Blood Pressure (mm Hg)					
Pre-Lisinopril value					
N	2	2	2	1	7
Mean (SD)	123.2 (7.8)	130.0 (4.2)	133.2 (9.7)	110.0	126.1 (9.8)
Median (Min, Max)	123.2 (117.7, 128.7)	130.0 (127.0, 133.0)	133.2 (126.3, 140.0)	110.0 (110.0, 110.0)	127.0 (110.0, 140.0)
Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	112.0 (29.7)	106.5 (3.5)	134.5 (14.8)	112.0	116.9 (18.4)
Median (Min, Max)	112.0 (91.0, 133.0)	106.5 (104.0, 109.0)	134.5 (124.0, 145.0)	112.0 (112.0, 112.0)	112.0 (91.0, 145.0)
Change from Baseline at Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	-11.2 (21.9)	-23.5 (7.8)	1.3 (5.2)	2.0	-9.2 (14.9)
Median (Min, Max)	-11.2 (-26.7, 4.3)	-23.5 (-29.0, -18.0)	1.3 (-2.3, 5.0)	2.0 (2.0, 2.0)	-2.3 (-29.0, 5.0)
95% CI for Median ¹					(-29.0, 5.0)
Percentage Change at Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	-9.7 (18.4)	-18.0 (5.4)	0.9 (3.8)	1.8	-7.4 (11.8)
Median (Min, Max)	-9.7 (-22.7, 3.4)	-18.0 (-21.8, -14.2)	0.9 (-1.8, 3.6)	1.8 (1.8, 1.8)	-1.9 (-22.7, 3.6)

Note: ¹ 95% Non-parametric Confidence Interval for median
Pre-lisinopril values are the average of the chart review records collected ≤6 weeks prior to first lisinopril dose recorded on prior and concomitant medications form.

Table 14.2.2.1.2.2
Summary of Diastolic Blood Pressure by Enrollment Group
PD Population - IC2 with Valid Pre-Lisinopril Records

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=10)
	0.1 mg/kg/day (N=2)	0.2 mg/kg/day (N=2)	0.1 mg/kg/day (N=5)	0.2 mg/kg/day (N=1)	
Diastolic Blood Pressure (mm Hg)					
Pre-Lisinopril value					
N	2	2	2	1	7
Mean (SD)	79.3 (0.5)	84.0 (9.9)	85.9 (12.1)	55.0	79.1 (12.7)
Median (Min, Max)	79.3 (79.0, 79.7)	84.0 (77.0, 91.0)	85.9 (77.3, 94.5)	55.0 (55.0, 55.0)	79.0 (55.0, 94.5)
Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	65.0 (1.4)	61.0 (11.3)	87.0 (19.8)	61.0	69.6 (15.2)
Median (Min, Max)	65.0 (64.0, 66.0)	61.0 (53.0, 69.0)	87.0 (73.0, 101.0)	61.0 (61.0, 61.0)	66.0 (53.0, 101.0)
Change from Baseline at Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	-14.3 (0.9)	-23.0 (1.4)	1.1 (7.7)	6.0	-9.5 (12.5)
Median (Min, Max)	-14.3 (-15.0, -13.7)	-23.0 (-24.0, -22.0)	1.1 (-4.3, 6.5)	6.0 (6.0, 6.0)	-13.7 (-24.0, 6.5)
95% CI for Median ¹					(-24.0, 6.5)
Percentage Change at Pre-Dose at PK Visit					
N	2	2	2	1	7
Mean (SD)	-18.1 (1.3)	-27.7 (4.9)	0.6 (8.8)	10.9	-11.3 (15.9)
Median (Min, Max)	-18.1 (-19.0, -17.2)	-27.7 (-31.2, -24.2)	0.6 (-5.6, 6.9)	10.9 (10.9, 10.9)	-17.2 (-31.2, 10.9)

Note:¹ 95% Non-parametric Confidence Interval for median
Pre-lisinopril values are the average of the chart review records collected <=6 weeks prior to first lisinopril dose recorded on prior and concomitant medications form.

Table 14.2.2.1.3.1
Systolic Blood Pressure Change from Baseline Estimated from Mixed Effects Model
PD Population - IC1

	Change from Baseline			Difference between 0.1 and 0.2 mg/kg ¹
	0.1 mg/kg	0.2 mg/kg	0.4 mg/kg	
Pre-Dose at PK Visit				
Estimate	-5.8	-8.3	-13	-2.5
95% Confidence Interval	(-14.4, 2.7)	(-16.9, 0.2)	(-26.8, 0.9)	(-14.6, 9.6)
10 Minutes Post-Iohexol				
Estimate	-11	-10.9	-29	0.1
95% Confidence Interval	(-19.6, -2.4)	(-19.9, -1.8)	(-42.8, -15.1)	(-12.3, 12.6)
4 Hours Post-Dose				
Estimate	-7.3	-14.5	-32.5	-7.1
95% Confidence Interval	(-15.9, 1.2)	(-23.5, -5.4)	(-46.3, -18.6)	(-19.6, 5.3)
8 Hours Post-Dose				
Estimate	-9.5	-17.9	-17	-8.4
95% Confidence Interval	(-18.1, -0.9)	(-26.9, -8.8)	(-30.8, -3.1)	(-20.8, 4.1)
12 Hours Post-Dose				
Estimate	-3.8	-8.5	-20	-4.6
95% Confidence Interval	(-12.4, 4.7)	(-17.5, 0.6)	(-33.8, -6.1)	(-17.1, 7.8)
24 Hours Post-Dose				
Estimate	-2.5	-14.5	-11	-12
95% Confidence Interval	(-11.1, 6.1)	(-23.5, -5.4)	(-24.8, 2.9)	(-24.4, 0.5)

Note: Change from baseline estimated as PK visit time point minus baseline model estimates.

¹ Estimated from 0.2 minus 0.1 mg/kg dose group model-estimated changes from baseline.

Parameters estimated from mixed effects model with subject random effect and time, dose group, and interaction fixed effects.

Table 14.2.2.1.3.2
Diastolic Blood Pressure Change from Baseline Estimated from Mixed Effects Model
PD Population - IC1

	Change from Baseline			Difference between 0.1 and 0.2 mg/kg ¹
	0.1 mg/kg	0.2 mg/kg	0.4 mg/kg	
Pre-Dose at PK Visit				
Estimate	-6.5	-3.8	-8.2	2.7
95% Confidence Interval	(-15.7, 2.7)	(-13, 5.3)	(-23.2, 6.8)	(-10.3, 15.7)
10 Minutes Post-Iohexol				
Estimate	-6.7	-11.3	-17.7	-4.7
95% Confidence Interval	(-15.8, 2.5)	(-21.1, -1.6)	(-32.7, -2.7)	(-18.1, 8.7)
4 Hours Post-Dose				
Estimate	-9.5	-14.9	-16.2	-5.4
95% Confidence Interval	(-18.7, -0.3)	(-24.7, -5.2)	(-31.2, -1.2)	(-18.8, 8)
8 Hours Post-Dose				
Estimate	-3.2	-19.9	-13.2	-16.8
95% Confidence Interval	(-12.3, 6)	(-29.7, -10.2)	(-28.2, 1.8)	(-30.2, -3.4)
12 Hours Post-Dose				
Estimate	-3.8	-9.3	-8.7	-5.5
95% Confidence Interval	(-13, 5.3)	(-19.1, 0.4)	(-23.7, 6.3)	(-18.9, 7.9)
24 Hours Post-Dose				
Estimate	-6	-14.9	-3.7	-8.9
95% Confidence Interval	(-15.2, 3.2)	(-24.7, -5.2)	(-18.7, 11.3)	(-22.3, 4.5)

Note: Change from baseline estimated as PK visit time point minus baseline model estimates.

¹ Estimated from 0.2 minus 0.1 mg/kg dose group model-estimated changes from baseline.

Parameters estimated from mixed effects model with subject random effect and time, dose group, and interaction fixed effects.

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Mean 24hr Systolic blood pressure (mm Hg)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	127.5 (4.9)	126.0	121.5 (7.8)	137.0 (0.0)	128.3 (7.5)
Median (Min, Max)	127.5 (124.0, 131.0)	126.0 (126.0, 126.0)	121.5 (116.0, 127.0)	137.0 (137.0, 137.0)	127.0 (116.0, 137.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	121.5 (9.2)	110.0	108.0 (4.2)	122.0 (8.5)	116.1 (8.9)
Median (Min, Max)	121.5 (115.0, 128.0)	110.0 (110.0, 110.0)	108.0 (105.0, 111.0)	122.0 (116.0, 128.0)	115.0 (105.0, 128.0)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-3.0	-16.0	-11.0	-15.0 (8.5)	-12.0 (6.9)
Median (Min, Max)	-3.0 (-3.0, -3.0)	-16.0 (-16.0, -16.0)	-11.0 (-11.0, -11.0)	-15.0 (-21.0, -9.0)	-11.0 (-21.0, -3.0)
95% CI for Median ¹					(-21.0, -3.0)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-2.3	-12.7	-9.5	-11.0 (6.2)	-9.3 (5.1)
Median (Min, Max)	-2.3 (-2.3, -2.3)	-12.7 (-12.7, -12.7)	-9.5 (-9.5, -9.5)	-11.0 (-15.3, -6.6)	-9.5 (-15.3, -2.3)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Mean Systolic daytime blood pressure (mm Hg)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	129.5 (3.5)	128.0	122.0 (7.1)	139.5 (0.7)	130.0 (7.9)
Median (Min, Max)	129.5 (127.0, 132.0)	128.0 (128.0, 128.0)	122.0 (117.0, 127.0)	139.5 (139.0, 140.0)	128.0 (117.0, 140.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	123.0 (9.9)	111.0	109.0 (7.1)	125.5 (7.8)	118.0 (9.8)
Median (Min, Max)	123.0 (116.0, 130.0)	111.0 (111.0, 111.0)	109.0 (104.0, 114.0)	125.5 (120.0, 131.0)	116.0 (104.0, 131.0)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-2.0	-17.0	-13.0	-14.0 (8.5)	-12.0 (7.2)
Median (Min, Max)	-2.0 (-2.0, -2.0)	-17.0 (-17.0, -17.0)	-13.0 (-13.0, -13.0)	-14.0 (-20.0, -8.0)	-13.0 (-20.0, -2.0)
95% CI for Median ¹					(-20.0, -2.0)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-1.5	-13.3	-11.1	-10.0 (6.0)	-9.2 (5.4)
Median (Min, Max)	-1.5 (-1.5, -1.5)	-13.3 (-13.3, -13.3)	-11.1 (-11.1, -11.1)	-10.0 (-14.3, -5.8)	-11.1 (-14.3, -1.5)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Mean Systolic nighttime blood pressure (mm Hg)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	121.0 (5.7)	118.0	119.5 (13.4)	126.0 (0.0)	121.6 (6.8)
Median (Min, Max)	121.0 (117.0, 125.0)	118.0 (118.0, 118.0)	119.5 (110.0, 129.0)	126.0 (126.0, 126.0)	125.0 (110.0, 129.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	111.0 (2.8)	102.0	107.5 (2.1)	112.5 (10.6)	109.1 (5.9)
Median (Min, Max)	111.0 (109.0, 113.0)	102.0 (102.0, 102.0)	107.5 (106.0, 109.0)	112.5 (105.0, 120.0)	109.0 (102.0, 120.0)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-12.0	-16.0	-4.0	-13.5 (10.6)	-11.8 (7.0)
Median (Min, Max)	-12.0 (-12.0, -12.0)	-16.0 (-16.0, -16.0)	-4.0 (-4.0, -4.0)	-13.5 (-21.0, -6.0)	-12.0 (-21.0, -4.0)
95% CI for Median ¹					(-21.0, -4.0)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-9.6	-13.6	-3.6	-10.7 (8.4)	-9.6 (5.6)
Median (Min, Max)	-9.6 (-9.6, -9.6)	-13.6 (-13.6, -13.6)	-3.6 (-3.6, -3.6)	-10.7 (-16.7, -4.8)	-9.6 (-16.7, -3.6)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
24hr Systolic blood pressure load (%)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	51.1 (13.2)	25.5	40.3 (7.8)	70.9 (12.9)	50.0 (18.5)
Median (Min, Max)	51.1 (41.8, 60.4)	25.5 (25.5, 25.5)	40.3 (34.8, 45.8)	70.9 (61.8, 80.0)	45.8 (25.5, 80.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	37.7 (18.7)	0.0	5.4 (3.0)	25.5 (6.4)	19.6 (17.8)
Median (Min, Max)	37.7 (24.4, 50.9)	0.0 (0.0, 0.0)	5.4 (3.3, 7.5)	25.5 (21.0, 30.0)	21.0 (0.0, 50.9)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-9.5	-25.5	-31.5	-45.4 (19.2)	-31.5 (17.9)
Median (Min, Max)	-9.5 (-9.5, -9.5)	-25.5 (-25.5, -25.5)	-31.5 (-31.5, -31.5)	-45.4 (-59.0, -31.8)	-31.5 (-59.0, -9.5)
95% CI for Median ¹					(-59.0, -9.5)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-15.7	-100.0	-90.5	-62.6 (15.8)	-66.3 (33.7)
Median (Min, Max)	-15.7 (-15.7, -15.7)	-100.0 (-100.0, -100.0)	-90.5 (-90.5, -90.5)	-62.6 (-73.8, -51.5)	-73.8 (-100.0, -15.7)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Nocturnal systolic dip (%)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	7.1 (2.3)	7.9	2.3 (5.3)	9.8 (0.5)	6.6 (3.9)
Median (Min, Max)	7.1 (5.4, 8.7)	7.9 (7.9, 7.9)	2.3 (-1.5, 6.0)	9.8 (9.4, 10.1)	7.9 (-1.5, 10.1)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	9.6 (4.9)	8.2	1.2 (4.4)	10.5 (3.0)	7.3 (5.1)
Median (Min, Max)	9.6 (6.1, 13.1)	8.2 (8.2, 8.2)	1.2 (-1.9, 4.3)	10.5 (8.4, 12.6)	8.2 (-1.9, 13.1)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	7.7	0.3	-7.9	0.8 (2.5)	0.3 (5.7)
Median (Min, Max)	7.7 (7.7, 7.7)	0.3 (0.3, 0.3)	-7.9 (-7.9, -7.9)	0.8 (-1.0, 2.5)	0.3 (-7.9, 7.7)
95% CI for Median ¹					(-7.9, 7.7)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	142.6	3.8	-131.7	7.1 (25.0)	5.8 (97.8)
Median (Min, Max)	142.6 (142.6, 142.6)	3.8 (3.8, 3.8)	-131.7 (-131.7, -131.7)	7.1 (-10.6, 24.8)	3.8 (-131.7, 142.6)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.1
Summary of Systolic ABPM and Overall Diagnosis by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Diagnosis					
Baseline					
Normal Blood Pressure (< 90th percentile)	0 (0.0%)	1 (100.0%)	0 (0.0%)	0 (0.0%)	1 (7.1%)
Pre-Hypertension (90- < 95th percentile)	1 (33.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (7.1%)
Stage 1 Hypertension (95th percentile- < 5mm Hg above 99th percentile)	1 (33.3%)	0 (0.0%)	2 (66.7%)	2 (40.0%)	5 (35.7%)
Stage 2 Hypertension (≥ 5mm Hg above 99th percentile)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Final Visit					
Normal Blood Pressure (< 90th percentile)	0 (0.0%)	1 (100.0%)	1 (33.3%)	1 (20.0%)	3 (21.4%)
Pre-Hypertension (90- < 95th percentile)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Stage 1 Hypertension (95th percentile- < 5mm Hg above 99th percentile)	2 (66.7%)	0 (0.0%)	1 (33.3%)	1 (20.0%)	4 (28.6%)
Stage 2 Hypertension (≥ 5mm Hg above 99th percentile)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Improvement from Baseline	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (20.0%)	1 (7.1%)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.2
Summary of Diastolic ABPM by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Mean 24hr Diastolic blood pressure (mm Hg)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	77.0 (2.8)	71.0	75.5 (4.9)	75.0 (5.7)	75.1 (3.8)
Median (Min, Max)	77.0 (75.0, 79.0)	71.0 (71.0, 71.0)	75.5 (72.0, 79.0)	75.0 (71.0, 79.0)	75.0 (71.0, 79.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	78.0 (5.7)	61.0	72.0 (5.7)	66.0 (5.7)	70.4 (7.6)
Median (Min, Max)	78.0 (74.0, 82.0)	61.0 (61.0, 61.0)	72.0 (68.0, 76.0)	66.0 (62.0, 70.0)	70.0 (61.0, 82.0)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-1.0	-10.0	-11.0	-9.0 (0.0)	-8.0 (4.0)
Median (Min, Max)	-1.0 (-1.0, -1.0)	-10.0 (-10.0, -10.0)	-11.0 (-11.0, -11.0)	-9.0 (-9.0, -9.0)	-9.0 (-11.0, -1.0)
95% CI for Median ¹					(-11.0, -1.0)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-1.3	-14.1	-13.9	-12.0 (0.9)	-10.7 (5.3)
Median (Min, Max)	-1.3 (-1.3, -1.3)	-14.1 (-14.1, -14.1)	-13.9 (-13.9, -13.9)	-12.0 (-12.7, -11.4)	-12.7 (-14.1, -1.3)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.2
Summary of Diastolic ABPM by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
24hr Diastolic blood pressure load (%)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	46.2 (9.3)	21.2	47.0 (13.4)	23.7 (7.6)	36.4 (14.7)
Median (Min, Max)	46.2 (39.6, 52.7)	21.2 (21.2, 21.2)	47.0 (37.5, 56.5)	23.7 (18.3, 29.1)	37.5 (18.3, 56.5)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	49.1 (30.6)	0.0	19.6 (6.9)	8.6 (4.8)	22.1 (23.5)
Median (Min, Max)	49.1 (27.4, 70.7)	0.0 (0.0, 0.0)	19.6 (14.7, 24.5)	8.6 (5.2, 12.0)	14.7 (0.0, 70.7)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	-12.2	-21.2	-41.8	-15.1 (2.8)	-21.1 (12.1)
Median (Min, Max)	-12.2 (-12.2, -12.2)	-21.2 (-21.2, -21.2)	-41.8 (-41.8, -41.8)	-15.1 (-17.1, -13.1)	-17.1 (-41.8, -12.2)
95% CI for Median ¹					(-41.8, -12.2)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	-30.8	-100.0	-74.0	-65.2 (9.1)	-67.0 (25.2)
Median (Min, Max)	-30.8 (-30.8, -30.8)	-100.0 (-100.0, -100.0)	-74.0 (-74.0, -74.0)	-65.2 (-71.6, -58.8)	-71.6 (-100.0, -30.8)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.2
Summary of Diastolic ABPM by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		>60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Mean Diastolic daytime blood pressure (mm Hg)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	78.5 (3.5)	72.0	76.0 (5.7)	77.0 (5.7)	76.4 (4.2)
Median (Min, Max)	78.5 (76.0, 81.0)	72.0 (72.0, 72.0)	76.0 (72.0, 80.0)	77.0 (73.0, 81.0)	76.0 (72.0, 81.0)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	80.5 (6.4)	62.0	73.0 (5.7)	68.0 (4.2)	72.1 (7.8)
Median (Min, Max)	80.5 (76.0, 85.0)	62.0 (62.0, 62.0)	73.0 (69.0, 77.0)	68.0 (65.0, 71.0)	71.0 (62.0, 85.0)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	0.0	-10.0	-11.0	-9.0 (1.4)	-7.8 (4.5)
Median (Min, Max)	0.0 (0.0, 0.0)	-10.0 (-10.0, -10.0)	-11.0 (-11.0, -11.0)	-9.0 (-10.0, -8.0)	-10.0 (-11.0, 0.0)
95% CI for Median ¹					(-11.0, 0.0)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	0.0	-13.9	-13.8	-11.7 (1.0)	-10.2 (5.8)
Median (Min, Max)	0.0 (0.0, 0.0)	-13.9 (-13.9, -13.9)	-13.8 (-13.8, -13.8)	-11.7 (-12.3, -11.0)	-12.4 (-13.9, 0.0)

Note:¹ 95% Non-parametric Confidence Interval for median

Table 14.2.2.2.2
Summary of Diastolic ABPM by Enrollment Group
PD Population - IC1

	30 – 59 ml/min per 1.73m ²		≥60 ml/min per 1.73m ²		Total (N=14)
	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=1)	0.1 mg/kg/day (N=3)	0.2 mg/kg/day (N=5)	
Nocturnal Diastolic dip (%)					
Pre-dose value					
N	2	1	2	2	7
Mean (SD)	11.0 (3.2)	9.8	3.2 (2.5)	12.2 (5.5)	8.9 (4.9)
Median (Min, Max)	11.0 (8.7, 13.2)	9.8 (9.8, 9.8)	3.2 (1.4, 5.0)	12.2 (8.3, 16.1)	8.7 (1.4, 16.1)
Post-dose value					
N	2	1	2	2	7
Mean (SD)	18.1 (0.6)	9.7	7.6 (4.6)	12.8 (8.1)	12.4 (5.9)
Median (Min, Max)	18.1 (17.7, 18.5)	9.7 (9.7, 9.7)	7.6 (4.3, 10.8)	12.8 (7.1, 18.5)	10.8 (4.3, 18.5)
Change from Baseline					
N	1	1	1	2	5
Mean (SD)	5.3	-0.1	-0.7	0.6 (13.6)	1.1 (7.2)
Median (Min, Max)	5.3 (5.3, 5.3)	-0.1 (-0.1, -0.1)	-0.7 (-0.7, -0.7)	0.6 (-9.0, 10.2)	-0.1 (-9.0, 10.2)
95% CI for Median ¹					(-9.0, 10.2)
Percentage Change					
N	1	1	1	2	5
Mean (SD)	40.2	-1.0	-14.0	33.5 (126.4)	18.4 (67.7)
Median (Min, Max)	40.2 (40.2, 40.2)	-1.0 (-1.0, -1.0)	-14.0 (-14.0, -14.0)	33.5 (-55.9, 122.9)	-1.0 (-55.9, 122.9)

Note:¹ 95% Non-parametric Confidence Interval for median