



Figure 14.1
Iohexol GFR (iGFR) and Estimated GFR (eGFR) Comparison
Safety Population



Note: CCC=0.69. Mean error=-11.16. Median % error=-14.42%.
Linear regression (solid) and unity (dotted) lines plotted.

Figure 14.2
Mean Blood Pressure Change Scores by Time
PD Population - IC1

Time Point	Systolic		Diastolic	
	0.1 mg/kg	0.2 mg/kg	0.1 mg/kg	0.2 mg/kg
Pre-Dose at PK Visit	-5.8±3.1	-8.3±4.3	-6.5±5.6	-3.8±3.3
10 Minutes Post-lohexol	-11±4.5	-13.2±3.4	-6.7±6.2	-11.2±4.3
4 Hours Post-Dose	-7.3±3.1	-16.8±5.7	-9.5±5.2	-14.8±3.2
8 Hours Post-Dose	-9.5±4.6	-20.2±4.9	-3.2±7.3	-19.8±5.3
12 Hours Post-Dose	-3.8±5.1	-10.8±4.7	-3.8±5.9	-9.2±2.7
24 Hours Post-Dose	-2.5±3	-16.8±4.9	-6±5.9	-14.8±2.9



Note: Systolic (left) and diastolic (right) BP for 0.1 (N=6) and 0.2 (N=6 at 0 hours; else N=5) mg/kg groups. Error bars are mean - 1 SE (plot) or ± 1 SE (table). Pre-dose values plotted at -30 minutes.